Appl. No.

: 10/063,713

Filed

.

May 8, 2002

AMENDMENTS TO THE CLAIMS

1-5. (Canceled).

- 6. (Previously Presented) An isolated nucleic acid comprising:
- (a) the nucleic acid sequence of SEQ ID NO:81;
- (b) the full-length coding sequence of the nucleic acid sequence of SEQ ID NO:81; or
- (c) the full-length coding sequence of the cDNA deposited under ATCC accession number 203317.

7-10. (Canceled).

- 11. (Previously Presented) The isolated nucleic acid of Claim 6 comprising the nucleic acid sequence of SEQ ID NO:81.
- 12. (Previously Presented) The isolated nucleic acid of Claim 6 comprising the full-length coding sequence of the nucleic acid sequence of SEQ ID NO:81.
- 13. (Original) The isolated nucleic acid of Claim 6 comprising the full-length coding sequence of the cDNA deposited under ATCC accession number 203317.
- 14. (Previously Presented) An isolated nucleic acid that hybridizes under stringent conditions to:
 - (a) the nucleic acid sequence of SEQ ID NO:81 or the complement thereof;
- (b) the full-length coding sequence of the nucleic acid sequence of SEQ ID NO:81 or the complement thereof; or
- (c) the full-length coding sequence of the cDNA deposited under ATCC accession number 203317 or the complement thereof;

wherein said stringent conditions comprise 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 μg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C;

wherein said isolated nucleic acid molecule is suitable for use as a PCR primer, or probe;

and wherein said isolated nucleic acid is at least about 450 nucleotides in length.

15. (Canceled).

OK to enter